

PhysiciansCARE®

ASPIRIN

Pain Reliever and Fever Reducer (NSAID)

COATED TABLETS

Tamper-Evident Packets of 2 Tablets
Do not use if packet is open or torn.

Aspirin (NSAID) 325 mg
100 Tablets (50 Packets, 2 tablets each)

Also Available from PhysiciansCare®
in individual dose packets:

PhysiciansCare® Non-Aspirin (Acetaminophen)

PhysiciansCare® Ibuprofen

PhysiciansCare® Cold & Cough

PhysiciansCare® Allergy

PhysiciansCare® Sinus

PhysiciansCare® Antacid

PhysiciansCARE®

ASPIRIN

Pain Reliever and Fever Reducer (NSAID)

COATED CAPLET

Individually packaged for workplace use

Tamper-Evident Packets of 2 Tablets
Do not use if packet is open or torn.

Aspirin (NSAID) 325 mg
100 Tablets (50 Packets, 2 tablets each)



Acme United Corporation

Legend

M-01
XL Dispenser (M Box)

Item 90014-005
Description Aspirin, 50 Packets

Version revG
Date 11.20 .23
Specs Refer to Dieline /
CMKY / w/Bleed

FAO/ANSI
Pantone & Process Colors

FAO Blue	ANSI Bandages	ANSI Antiseptics
CMYK 100/70/0/0 PMS286C	CMYK 0/18/100/0 PMS123C	CMYK 100/75/0/0 PMS287C
ANSI Burn Treatment	ANSI Personal Protection Equipment	ANSI Misc
CMYK 0/100/76/5 PMS186C	CMYK 0/56/87/0 PMS144C	CMYK 100/0/69/43 PMS342C

Rev. 8-11-23
Mfg. 24

Drug Facts provided by Acme

Retain carton for complete product information

Drug Facts

Active ingredient (in each tablet) Aspirin (NSAID)* 325 mg
Purpose Pain reliever / fever reducer
*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves:

- headache
- minor arthritis pain
- pain & fever of colds
- cramps with pain
- toothache
- aches with pain

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcohol drinks every day while using this product
■ take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have asthma
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Drug Facts (continued)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- if ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose

Adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours

Children under 12 years: consult a doctor

Other information

- read all product information before using
- store at room temperature
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions? 1-800-835-2263

Contact Us 1-800-835-2263 www.FirstAidOnly.com

VOID AQUEOUS ZONE
Does Not Print!

Manufactured for:
Acme United Corporation
1 Waterview Dr. Shelton, CT 06484
www.FirstAidOnly.com
© 2023 Acme United Corporation.

217914-revG



0 73577 90014 1