

McNEIL P/N 8737970
CTN MOTRIN IB CPLT 100 DR

8737970



Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk, if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Important: Read all product information before using. Keep this box for important information.

Drug Facts

Active ingredient (in each caplet) Purpose
Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (Wheezing)
- shock
- skin redness
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- use other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

8737970

EXP: 3 0045-0481-016



NDC 50580-110-10

Motrin[®] IB

Ibuprofen Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)

100 Coated Caplets*

*Capsule-Shaped Tablets

Drug Facts (continued)

polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide

Questions or comments?
Call 1-877-895-3665: weekdays 8:00 AM to 8:00 PM EST

McNeil
Distributed by:
McNeil Consumer Healthcare
DIVISION OF MCNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
©MCN-PPC, INC. 2011
www.motrin.com

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older	take 1 caplet every 4 to 6 hours while symptoms persist
children under 12 years	if pain or fever does not respond to 1 caplet, 2 caplets may be used
	do not exceed 6 caplets in 24 hours, unless directed by a doctor

ask a doctor

Other information

- store between 20-25°C (68-77°F)
- do not use if neck wrap or foil inner seal imprinted "SAFETY SEAL®" is broken or missing
- see end panel for lot number and expiration date

Inactive ingredients cambeaba wax, colloidal silicon dioxide, corn starch, FD&C yellow no. 6, hypromellose, iron oxide, magnesium stearate, polyethylene glycol, polyvinylpyrrolidone



08737970

ORG YELLOW BLUE BLK

APPROVED FOR RELEASE TO PRODUCTION

by: **GREAT ATLANTIC** as authorized by: **McNeil**

Approver #1

Signature: Ashley Keisling

Name: Ashley Keisling

Title: Production Coordinator

Date: 0)!")!\$" ##

Approver #2

Signature: Diane O'Dwyer

Name: Diane O'Dwyer

Title: Packaging Graphics Manager

Date: 0)!")!\$" ##

Copy & content compared to original document no. >55 \$" ##Z' %#"