



**Acme United Corporation** 

Legend

# M-01 **XL Dispenser (M Box)**

Allergy Plus, 50 Packets 10.20.17

Refer to Dieline / CMKY / w/Bleed FAO/ANSI Pantone & Process Colors

90091

Rev. 10/20/17 Mfg. 1, 14

90091

### Medique

# Also Available from **Physicians** *Care*® in individual dose packets:

Physicians Care® Aspirin

**Physicians** *Care* Non-Aspirin (Acetaminophen)

Physicians Care® Cold & Cough

Physicians Care® Allergy

**Physicians** Care® Sinus

Physicians Care® Antacid

Physicians Care® Extra Strength Pain Relief

**Physicians** Care® Motion Sickness



**Multi-Symptom Relief** 



THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

COATED TABLETS

Tamper-Evident Packets of 2 Tablets

100 Tablets (50 Packets, 2 tablets each

Retain carton for complete product information

# Drug Facts

runny nose and sneezingminor aches and pains

sinus congestion and pressur itching of the nose or throat

Liver warning: This product contains acetaminophen.
Severe liver damage may occur if you take:

more than 12 tablets in 24 hours, which is the maximum daily amo

with other drugs containing acetaminophen
 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: skin reddening blisters rash

If a skin reaction occurs, stop use and seek medical help right away.

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

for more than 10 days for pain unless directed by a doctor ■ for more than 10 days for fever unless directed by a doctor
■ with any other drug using the above active ingredients

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an

Ask a doctor before use if you have
■ liver disease ■ thyroid disease ■ diabetes ■ high blood pressure
■ heart disease ■ glaucoma ■ chronic bronchitis or emphysema difficulty in urination due to enlargement of the prostate gland

# If pregnant or breast-feeding, ask a health profess

Drug Facts (continued)

Stop use and ask a doctor if

any new symptoms appear redness or swelling is present

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

Do not exceed recommended dosage. If nervousness, dizziness or sleeplessness occurs, discontinue use and consult a docto

Ask a doctor or pharmacist before use if you are
 taking the blood thinning drug warfarin
 taking sedatives or tranquilizers

supplemental set a doctor if
symptoms do not improve within 7 days or are accompanied by a fever
pain or fever persists or gets worse

# Directions ■ do not use more that

Adults and children: (12 years and older) Take 2 tablets every 4 hours. Do not take more than 12 tablets in 24 hours.

Children under

Other information
■ store at room temperature 59°-86°F (15°-30°C)
■ avoid excessive heat and humidity
■ tamper evident sealed packets

# do not use any opened or torn pack

Inactive ingredients
corn starch, crospovidone\*, hypromellose, maltodextrin\*,
microcrystalline cellulose, polyethylene glycol, povidone,
pregelatinized starch\*, sodium starch glycolate\*, stearic acid

▶ **Questions?** 1-800-835-2263

Contact Us 1-800-835-2263 www.FirstAidOnly.com





**Multi-Symptom Relief** 



THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

COATED TABLETS

Tamper-Evident Packets of 2 Tablets

100 Tablets (50 Packets, 2 tablets each





217991-revF